

Correspondence Information

Correspondence Customer	22847
-------------------------	-------

Number::

Representative Information

Representative Customer Number::	22847
----------------------------------	-------

[This application has no priority claims or assignee data]

[illegible]

INBRED MAIZE LINE NP2222

FIELD OF THE INVENTION

This invention is in the field of maize breeding, specifically relating to an inbred
5 maize line designated NP2222.

BACKGROUND OF THE INVENTION

The goal of plant breeding is to combine in a single variety or hybrid various
desirable traits. For field crops, these traits may include resistance to diseases and insects,
10 tolerance to heat and drought, reducing the time to crop maturity, greater yield, and better
agronomic quality. With mechanical harvesting of many crops, uniformity of plant
characteristics such as germination and stand establishment, growth rate, maturity, and
plant and ear height, is important.

Field crops are bred through techniques that take advantage of the plant's method
15 of pollination. A plant is self-pollinated if pollen from one flower is transferred to the
same or another flower of the same plant. A plant is cross-pollinated if the pollen comes
from a flower on a different plant. Plants that have been self-pollinated and selected for
type for many generations become homozygous at almost all gene loci and produce a
uniform population of true breeding progeny. A cross between two different homozygous
20 lines produces a uniform population of hybrid plants that may be heterozygous for many
gene loci. A cross of two plants each heterozygous at a number of gene loci will produce
a population of hybrid plants that differ genetically and will not be uniform.

Maize (*Zea mays* L.), often referred to as corn in the United States, can be bred by
both self-pollination and cross-pollination techniques. Maize has separate male and
25 female flowers on the same plant, located on the tassel and the ear, respectively. Natural
pollination occurs in maize when wind blows pollen from the tassels to the silks that
protrude from the tops of the ears.

A reliable method of controlling male fertility in plants offers the opportunity for
improved plant breeding. This is especially true for development of maize hybrids, which
30 relies upon some sort of male sterility system. There are several options for controlling

male fertility available to breeders, such as: manual or mechanical emasculation (or detasseling), cytoplasmic male sterility, genetic male sterility, gametocides and the like.

Hybrid maize seed is typically produced by a male sterility system incorporating manual or mechanical detasseling. Alternate strips of two maize inbreds are planted in a field, and the pollen-bearing tassels are removed from one of the inbreds (female). Providing that there is sufficient isolation from sources of foreign maize pollen, the ears of the detasseled inbred will be fertilized only from the other inbred (male) and the resulting seed is therefore hybrid and will form hybrid plants.

The laborious, and occasionally unreliable, detasseling process can be avoided by using cytoplasmic male-sterile (CMS) inbreds. Plants of a CMS inbred are male sterile as a result of factors resulting from the cytoplasmic, as opposed to the nuclear, genome. Thus, this characteristic is inherited exclusively through the female parent in maize plants, since only the female provides cytoplasm to the fertilized seed. CMS plants are fertilized with pollen from another inbred that is not male-sterile. Pollen from the second inbred may or may not contribute genes that make the hybrid plants male-fertile. Seed from detasseled fertile maize and CMS produced seed of the same hybrid can be blended to insure that adequate pollen loads are available for fertilization when the hybrid plants are grown.

There are several methods of conferring genetic male sterility available, such as multiple mutant genes at separate locations within the genome that confer male sterility, as disclosed in U.S. Pat. Nos. 4,654,465 and 4,727,219 and chromosomal translocations as described in U.S. Pat. Nos. 3,861,709 and 3,710,511, the disclosures of which are specifically incorporated herein by reference. There are many other methods of conferring genetic male sterility in the art, each with its own benefits and drawbacks. These methods use a variety of approaches such as delivering into the plant a gene encoding a cytotoxic substance associated with a male tissue specific promoter or an antisense system in which a gene critical to fertility is identified and an antisense to that gene is inserted in the plant (EPO 89/3010153.8 and WO 90/08828).

Another system useful in controlling male sterility makes use of gametocides. Gametocides are not a genetic system, but rather a topical application of chemicals. These chemicals affect cells that are critical to male fertility. The application of these

chemicals affects fertility in the plants only for the growing season in which the gametocide is applied (see Carlson, Glenn R., U.S. Pat. No. 4,936,904, which is incorporated herein by reference). Application of the gametocide, timing of the application and genotype specificity often limit the usefulness of the approach.

5 The use of male sterile inbreds is but one factor in the production of maize hybrids. The development of maize hybrids requires, in general, the development of homozygous inbred lines, the crossing of these lines, and the evaluation of the crosses. Pedigree breeding and recurrent selection breeding methods are used to develop inbred lines from breeding populations. Breeding programs combine the genetic backgrounds
10 from two or more inbred lines or various other germplasm sources into breeding pools from which new inbred lines are developed by selfing and selection of desired phenotypes. The new inbreds are crossed with other inbred lines and the hybrids from these crosses are evaluated to determine which of those have commercial potential. Plant breeding and hybrid development are expensive and time-consuming processes.

15 Pedigree breeding starts with the crossing of two genotypes, each of which may have one or more desirable characteristics that is lacking in the other or which complements the other. If the two original parents do not provide all the desired characteristics, other sources can be included in the breeding population. In the pedigree method, superior plants are selfed and selected in successive generations. In the
20 succeeding generations the heterozygous condition gives way to homogeneous lines as a result of self-pollination and selection. Typically in the pedigree method of breeding five or more generations of selfing and selection is practiced: F1 to F2; F3 to F4; F4 to F5, etc.

25 Recurrent selection breeding, backcrossing for example, can be used to improve populations of either self or cross-pollinating crops. Backcrossing can be used to transfer a specific desirable trait from one inbred or source to an inbred that lacks the trait. This can be accomplished, for example, by first a superior inbred (recurrent parent) to a donor inbred (non-recurrent parent), that carries the appropriate gene(s) for the trait in question. The progeny of this cross is then mated back to the superior recurrent parent followed by
30 selection in the resultant progeny for the desired trait to be transferred from the non-recurrent parent. After five or more backcross generations with selection for the desired

trait, the progeny will be homozygous for loci controlling the characteristic being transferred, but will be like the superior parent for essentially all other genes. The last backcross generation is, then selfed to give pure breeding progeny for the gene(s) being transferred. A hybrid developed from inbreds containing the transferred gene(s) is essentially the same as a hybrid developed from the same inbreds without the transferred genes. As the varieties developed using recurrent selection breeding contain almost all of the characteristics of the recurrent parent, selecting a superior recurrent parent is desirable.

A single cross maize hybrid results from the cross of two inbred lines, each of which has a genotype that complements the genotype of the other. The hybrid progeny of the first generation is designated F1. In the development of commercial hybrids only the F1 hybrid plants are sought. Preferred F1 hybrids are more vigorous than their inbred parents. This hybrid vigor, or heterosis, can be manifested in many polygenic traits, including increased vegetative growth and increased yield.

The development of a maize hybrid involves three steps: (1) the selection of plants from various germplasm pools for initial breeding crosses; (2) the selfing of the selected plants from the breeding crosses for several generations to produce a series of inbred lines, which, although different from each other, breed true and are highly uniform; and (3) crossing the selected inbred lines with different inbred lines to produce the hybrid progeny (F1). During the inbreeding process in maize, the vigor of the lines decreases. Vigor is restored when two different inbred lines are crossed to produce the hybrid progeny (F1). An important consequence of the homozygosity and homogeneity of the inbred lines is that the hybrid between a defined pair of inbreds will always be the same. Once the inbreds that give a superior hybrid have been identified, the hybrid seed can be reproduced indefinitely as long as the homogeneity of the inbred parents is maintained.

A single cross hybrid is produced when two inbred lines are crossed to produce the F1 progeny. A double cross hybrid is produced from four inbred lines crossed in pairs ($A \times B$ and $C \times D$) and then the two F1 hybrids are crossed again $(A \times B) \times (C \times D)$. Much of the hybrid vigor exhibited by F1 hybrids is lost in the next generation (F2). Consequently, seed from hybrids is not used for planting stock.

Hybrid seed production requires elimination or inactivation of pollen produced by the female parent. Incomplete removal or inactivation of the pollen provides the potential for self-pollination. This inadvertently self-pollinated seed may be unintentionally harvested and packaged with hybrid seed. Once the seed is planted, it is possible to
5 identify and select these self-pollinated plants. These self-pollinated plants will be genetically equivalent to the female inbred line used to produce the hybrid. Typically these self-pollinated plants can be identified and selected due to their decreased vigor. Female selfs are identified by their less vigorous appearance for vegetative and/or reproductive characteristics, including shorter plant height, small ear size, ear and kernel
10 shape, cob color, or other characteristics.

Identification of these self-pollinated lines can also be accomplished through molecular marker analyses. See, "The Identification of Female Selfs in Hybrid Maize: A Comparison Using Electrophoresis and Morphology", Smith, J. S. C. and Wych, R. D., Seed Science and Technology 14, pp. 1-8 (1995), the disclosure of which is expressly
15 incorporated herein by reference. Through these technologies, the homozygosity of the self-pollinated line can be verified by analyzing allelic composition at various loci along the genome. Those methods allow for rapid identification of the invention disclosed herein. See also, "Identification of Atypical Plants in Hybrid Maize Seed by Postcontrol and Electrophoresis" Sarca, V. et al., Probleme de Genetica Teoritca si Aplicata Vol. 20
20 (1) p. 29-42.

As is readily apparent to one skilled in the art, the foregoing describes only two of the various ways by which the inbred can be obtained by those looking to use the germplasm. Other means are available, and the above examples are illustrative only.

Maize is an important and valuable field crop. Thus, a continuing goal of plant
25 breeders is to develop high-yielding maize hybrids that are agronomically sound based on stable inbred lines. The reasons for this goal are obvious: to maximize the amount of grain produced with the inputs used and minimize susceptibility of the crop to pests and environmental stresses. To accomplish this goal, the maize breeder must select and develop superior inbred parental lines for producing hybrids. This requires identification
30 and selection of genetically unique individuals that occur in a segregating population. The segregating population is the result of a combination of crossover events plus the

independent assortment of specific combinations of alleles at many gene loci that results in specific genotypes. The probability of selecting any one individual with a specific genotype from a breeding cross is infinitesimal due to the large number of segregating genes and the unlimited recombinations of these genes, some of which may be closely
5 linked. However, the genetic variation among individual progeny of a breeding cross allows for the identification of rare and valuable new genotypes. These new genotypes are neither predictable nor incremental in value, but rather the result of manifested genetic variation combined with selection methods, environments and the actions of the breeder. Thus, even if the entire genotypes of the parents of the breeding cross were
10 characterized and a desired genotype known, only a few, if any, individuals having the desired genotype may be found in a large segregating F2 population. Typically, however, neither the genotypes of the breeding cross parents nor the desired genotype to be selected is known in any detail. In addition, it is not known how the desired genotype would react with the environment. This genotype by environment interaction is an
15 important, yet unpredictable, factor in plant breeding. A breeder of ordinary skill in the art cannot predict the genotype, how that genotype will interact with various climatic conditions or the resulting phenotypes of the developing lines, except perhaps in a very broad and general fashion. A breeder of ordinary skill in the art would also be unable to recreate the same line twice from the very same original parents, as the breeder is unable
20 to direct how the genomes combine or how they will interact with the environmental conditions. This unpredictability results in the expenditure of large amounts of research resources in the development of a superior new maize inbred line.

SUMMARY OF THE INVENTION

25 According to the invention, there is provided a novel inbred maize line, designated NP2222. This invention thus relates to the seeds of inbred maize line NP2222, to the plants of inbred maize line NP2222, and to methods for producing a maize plant by crossing the inbred line NP2222 with itself or another maize line. This invention further relates to hybrid maize seeds and plants produced by crossing the inbred line NP2222
30 with another maize line. The invention further relates to methods for producing maize

lines derived from inbred maize line NP2222 and from hybrid maize lines developed therefrom, and to maize lines derived by use of those methods.

The invention is also directed to inbred maize line NP2222 into which one or more specific, single gene traits, for example transgenes, have been introgressed from another maize line. Preferably, the resulting line has essentially all of the morphological and physiological characteristics of inbred maize line of NP2222, in addition to the one or more specific, single gene traits introgressed into the inbred, preferably the resulting line has all of the morphological and physiological characteristics of inbred maize line of NP2222, in addition to the one or more specific, single gene traits introgressed into the inbred. The invention also relates to seeds of an inbred maize line NP2222 into which one or more specific, single gene traits have been introgressed and to plants of an inbred maize line NP2222 into which one or more specific, single gene traits have been introgressed. The invention further relates to methods for producing a maize plant by crossing plants of an inbred maize line NP2222 into which one or more specific, single gene traits have been introgressed with themselves or with another maize line. The invention also further relates to hybrid maize seeds and plants produced by crossing plants of an inbred maize line NP2222 into which one or more specific, single gene traits have been introgressed with another maize line. The invention is also directed to a method of producing inbreds comprising planting a collection of hybrid seed, growing plants from the collection, identifying inbreds among the hybrid plants, selecting the inbred plants and controlling their pollination to preserve their homozygosity.

DETAILED DESCRIPTION OF THE INVENTION

Inbred maize lines are typically developed for use in the production of hybrid maize lines. Inbred maize lines need to be highly homogeneous, homozygous and reproducible to be useful as parents of commercial hybrids. There are many analytical methods available to determine the homozygotic and phenotypic stability of these inbred lines.

The oldest and most traditional method of analysis is the observation of phenotypic traits. The data is usually collected in field experiments over the life of the maize plants to be examined. Phenotypic characteristics most often observed are for traits

associated with plant morphology, ear and kernel morphology, insect and disease resistance, maturity, and yield.

In addition to phenotypic observations, the genotype of a plant can also be examined. There are many laboratory-based techniques available for the analysis, comparison and characterization of plant genotype; among these are Isozyme Electrophoresis, Restriction Fragment Length Polymorphisms (RFLPs), Randomly Amplified Polymorphic DNAs (RAPDs), Arbitrarily Primed Polymerase Chain Reaction (AP-PCR), DNA Amplification Fingerprinting (DAF), Sequence Characterized Amplified Regions (SCARs), Amplified Fragment Length Polymorphisms (AFLPs), and Simple Sequence Repeats (SSRs) which are also referred to as Microsatellites.

Some of the most widely used of these laboratory techniques are Isozyme Electrophoresis and RFLPs as discussed in Lee, M., "Inbred Lines of Maize and Their Molecular Markers," The Maize Handbook, (Springer-Verlag, New York, Inc. 1994, at 423-432). Isozyme Electrophoresis is a useful tool in determining genetic composition, although it has relatively low number of available markers and the low number of allelic variants among maize inbreds. RFLPs have the advantage of revealing an exceptionally high degree of allelic variation in maize and the number of available markers is almost limitless. Maize RFLP linkage maps have been rapidly constructed and widely implemented in genetic studies. One such study is described in Boppenmaier, et al., "Comparisons among strains of inbreds for RFLPs", Maize Genetics Cooperative Newsletter, 65:1991, pg. 90. This study used 101 RFLP markers to analyze the patterns of 2 to 3 different deposits each of five different inbred lines. The inbred lines had been selfed from 9 to 12 times before being adopted into 2 to 3 different breeding programs. It was results from these 2 to 3 different breeding programs that supplied the different deposits for analysis. These five lines were maintained in the separate breeding programs by selfing or sibbing and rogueing off-type plants for an additional one to eight generations. After the RFLP analysis was completed, it was determined the five lines showed 0-2% residual heterozygosity. Although this was a relatively small study, it can be seen using RFLPs that the lines had been highly homozygous prior to the separate strain maintenance.

The production of hybrid maize lines typically comprises planting in pollinating proximity seeds of, for example, inbred maize line NP2222 and of a different inbred parent maize plant, cultivating the seeds of inbred maize line NP2222 and of said different inbred parent maize plant into plants that bear flowers, emasculating the male flowers of inbred maize line NP2222 or the male flowers of said different inbred parent maize plant to produce an emasculated maize plant, allowing cross-pollination to occur between inbred maize line NP2222 and said different inbred parent maize plant and harvesting seeds produced on said emasculated maize plant. The harvested seed are grown to produce hybrid maize plants.

Inbred maize line NP2222 can be crossed to inbred maize lines of various heterotic group (see e.g. Hallauer et al. (1988) in Corn and Corn Improvement, Sprague et al, eds, chapter 8, pages 463-564) for the production of hybrid maize lines.

TABLE I
VARIETY DESCRIPTION INFORMATION

Inbred maize line NP2222 is compared to inbred A632

	NP2222		A632	
Plant data	Mean	std. dev	Mean	std dev
Plant height (cm) (to tassel tip)	218.7	19.0	216.8	16.5
Ear Height (cm) (to base of top ear node)	072.0	9.9	093.2	9.0
Length of Top Ear Internode (cm)	015.7	0.8	014.1	1.5
Number of tillers	0.0	0.0	0.0	0.1
Ears per stalk	1.1	0.2	1.2	0.3
Leaf data	Mean	std. dev.	Mean	std dev
Width of ear node leaf (cm)	009.3	0.4	008.4	0.3
Length of ear node leaf (cm)	075.8	3.6	081.7	7.5
Number of leaves above top ear	06	0.4	06	0.3
Leaf angle (measure from 2 nd leaf above ear at Anthesis to stalk above leaf)	044	29.1	063	15.4
Tassel data	Mean	std dev	Mean	std dev
Number of primary lateral branches	06	0.7	08	1.2
Branch angle (degrees from central spike)	017	4.9	044	2.9

Tassel length (cm) (from top leaf collar to tassel tip)	47.2	2.4	35.3	2.5
Ear data (husked data)	Mean	std dev	Mean	std dev
Ear length (cm)	16.9	1.4	15.5	1.2
Ear diameter at midpoint (mm)	39.6	1.6	37.0	0.9
Ear weight (grams)	128.4	31.6	077.8	6.4
Number of Kernel Rows	15	0.7	16	0.6
Kernel length (mm) (dried)	11.5	1.1	09.5	0.4
Kernel width (mm) (dried)	07.4	0.2	07.1	0.7
Kernel thickness (mm) (dried)	3.6	0.6	04.4	0.4
Percentage of round kernels (shape grade) (dried)	4.1	2.7	46.8	8.2
Weight of 100 kernels (grams) (dried)	21.9		22.1	4.2
Cob diameter at midpoint (mm)	21.3	1.8	22.1	4.2
Descriptive ratings	NP2222	A632		
Leaf sheath pubescence (1=none to 9=many)	2	6		
Marginal waves (1=none to 9=many)	6	5		
Longitudinal creases (1=none to 9=many)	5	6		
Pollen shed (0= male sterile to 9=heavy shed)	6	6		
Position of ear at dry husk stage (1=upright, 2=horizontal, 3=pendent) (unhusked data)	1	2		
Husk tightness (1=very loose to 9=very tight) (unhusked data)	5	6		
Husk extension (1=short, ears exposed; 2=medium, <8cm; 3=long, 8-10 cm beyond ear tip; 4=very long, >10cm) (unhusked data)	2	3		
Kernel rows (1=indistinct, 2=distinct)	2	2		
Row alignment (1=straight, 2=slightly curved, 3=spiral)	1	1		
Ear taper (1=slight, 2=average, 3=extreme)	2	2		
Aleurone color pattern (1=homozygous, 2=segregating)	1	1		
Endosperm type (1=sweet, su1; 2=extra sweet, sh2; 3=normal)	3	3		

starch; 4=highamylose starch; 5=waxy starch; 6=high protein; 7=high lysine; 8=supersweet, se; 9=high oil; 10= other				
Anthocyanin of brace roots (1=absent, 2=faint, 3=moderate, 4=dark)	4	4		
Bar Glumes (Glume bands) 1=absent, 2=present	2	2		
	NP2222		A632	
Maturity	Days	Heat Units	Days	Heat Units
Emergence of 50% plants in silk	067	1247.7	072	1360.3
Emergence of 50% plants in pollen	066	1240.1	070	1310.7
From 50% silk to harvest at 25% moisture		1400		1384
From 10% to 90% pollen shed	003	0068.2	003	0091.2
Color data	NP2222		A632	
	Munsell (M)		Munsell	
Leaf	02	(5GY 4/4)	03	(5GY 3/4)
Anther	05	(2.5GY 8/8)	26	
Glume	26		26	
Silk (3 days after emergence) (unhusked data)	26		05	(2.5GY 8/6)
Fresh husk (25 days after 50% silking) (unhusked data)	05	(2.5GY 8/6)	05	(2.5GY 8/8)
Dry husk (65 days after 50% silking) (unhusked data)	22	(2.5Y 8/4)	22	(2.5Y 8/4)
Aleurone	18		26	
Cob	14	(5R 4/8)	14	(5R 4/8)
Hard endosperm	06	(2.5Y 8/10)	06	(.5Y 8/10)
Disease Resistance				
	NP2222		A632	
Common Rust	4		6	
Eyespot	5		6	
Northern Leaf Blight (Race mixed 1 & 2)	6		7	
B. Insect Resistance				
1 st generation (typically whorl leaf feeding)	2		4	
Agronomic Traits				
% Dropped Ears (at 65 days after anthesis)	0		5	

% Pre-anthesis Brittle snapping	1	0
% Pre-anthesis Root Lodging	1	3
% Post-anthesis Root Lodging	0	3
Kg/ha Yeild of Inbred Per Se (at 12-13% grain moisture)	5537.3	2194.9

All linear measurements are in centimeters unless otherwise noted.

In interpreting the foregoing color designations, reference may be made to the
 5 Munsell Glossy Book of Color, a standard color reference. Color codes: 1. light green,
 2. medium green, 3. dark green, 4. very dark green, 5. green-yellow, 6. pale yellow,
 7. yellow, 8. yellow-orange, 9. salmon, 10. pink-orange, 11. pink 12. light red, 13.
 cherry red, 14. red, 15. red and white, 16. pale purple, 17. purple, 18. colorless, 19.
 white, 20. white capped, 21. buff, 22. tan, 23. brown, 24. bronze, 25. variegated,
 10 26. other. Heat Units per day were calculated using the standard formula:
 $HU = \{MaxTemp (86) + Min Temp (50)\} / 2 - 50$.

NP2222 differs from A632 for several different traits. These traits are:

NP2222 is an earlier maturity inbred than A632. The heat unit accumulation to
 15 all of the flowering stages is less for NP2222 than A632. The heat units to 10% Pollen
 Shed is 1213 for NP2222 and is 1280 for A632. Heat units to 50% Pollen Shed is 1240
 for NP2222 as compared to 1311 for A632. Heat units to 50% Silk Emergence is 1248
 for NP2222 and 1360 for A632.

The days to 50% silk emergence is also significantly different for NP2222 than
 20 A632. NP2222 reaches 50% silk emergence in 67 days maturity as compared to 72 days
 for A632.

NP2222 has a significantly lower ear placement than A632. The ear height of
 NP2222 is 72 cm. The ear height of A632 is 93 cm.

The Length of the Top Ear Internode is significantly longer on NP2222 at 16 cm
 25 as compared to 14 on A632.

The NP2222 leaf color is lighter than the A632 leaf. The NP2222 leaf is a
 medium green (02 or 5GY 4/4) and the A632 leaf is dark green (03 or 5GY 3/4).

The Leaf Sheath Pubescence on NP2222 is rated a 2 and is significantly different
 than A632 rated a 6.

Some of the most distinct differences of NP2222 and A632 occur in the tassel. The most pronounced of these differences is the Tassel Branch Angle. The Tassel Branch Angle of NP2222 is 17 degrees while the branch angle for A632 is 44 degrees. NP2222 has a very upright tassel branch angle. The tassel length of NP2222 is also significantly longer than A632. The NP2222 Tassel is 47 cm long while the A632 Tassel is 35 cm.

The tassels of these two inbreds also differ in coloration. The anther color of the two inbreds is significantly different. The NP2222 anther color is green-yellow (05 or 2.5GY 8/6). The A632 anther color is green yellow with pale purple shading. The glume color of NP2222 is green-yellow (05 or 2.5GY 8/6) with purple shading. The glume color of A632 is is green-yellow (05 or 5GY 5/6) with the purple coloration (only) on the glume margins. The glume color bars of A632 have a slight purple shade. The glume color bars of NP2222 are light green to green-yellow.

Another pronounced color difference of these two inbreds is the color of the silk. The silk color of NP2222 is green-yellow (05 or 2.5GY 8/6) with purple shaded ends. The purple shaded ends of NP2222 are 17 or 5RP 3/8. The A632 silk is green-yellow (05 or 2.5GY 8/6).

The ear of NP2222 is different than the ear of A632. The Ear Diameter of NP2222 is 40 cm while the ear of A632 is 37 cm. The Ear Weight of NP2222 is 128 gm as compared to 78 gm of A632. The Cob diameter of NP2222 is significantly smaller at 21 mm than A632 at 23 mm. NP2222 has 4 percent Round Kernels on the ear as compared to 23 percent on A632. The NP2222 plant has fewer dropped ears at harvest than A632. The Dropped Ear Percentage for NP2222 is 0 while A632 is 5.

The invention also encompasses plants of inbred maize line NP2222 and parts thereof further comprising one or more specific, single gene traits which have been introgressed into inbred maize line NP2222 from another maize line. Preferably, one or more new traits are transferred to inbred maize line NP2222, or, alternatively, one or more traits of inbred maize line NP2222 are altered or substituted. The transfer (or introgression) of the trait(s) into inbred maize line NP2222 is for example achieved by recurrent selection breeding, for example by backcrossing. In this case, inbred maize line NP2222 (the recurrent parent) is first crossed to a donor inbred (the non-recurrent parent)

that carries the appropriate gene(s) for the trait(s) in question. The progeny of this cross is then mated back to the recurrent parent followed by selection in the resultant progeny for the desired trait(s) to be transferred from the non-recurrent parent. After three, preferably four, more preferably five or more generations of backcrosses with the recurrent parent with selection for the desired trait(s), the progeny will be heterozygous for loci controlling the trait(s) being transferred, but will be like the recurrent parent for most or almost all other genes (see, for example, Poehlman & Sleper (1995) *Breeding Field Crops*, 4th Ed., 172-175; Fehr (1987) *Principles of Cultivar Development*, Vol. 1: Theory and Technique, 360-376).

The laboratory-based techniques described above, in particular RFLP and SSR, are routinely used in such backcrosses to identify the progenies having the highest degree of genetic identity with the recurrent parent. This permits to accelerate the production of inbred maize lines having at least 90%, preferably at least 95%, more preferably at least 99% genetic identity with the recurrent parent, yet more preferably genetically identical to the recurrent parent, and further comprising the trait(s) introgressed from the donor parent. Such determination of genetic identity is based on molecular markers used in the laboratory-based techniques described above. Such molecular markers are for example those known in the art and described in Boppenmaier, et al., "Comparisons among strains of inbreds for RFLPs", *Maize Genetics Cooperative Newsletter* (1991) 65, pg. 90, or those available from the University of Missouri database and the Brookhaven laboratory database (see <http://www.agron.missouri.edu>). The last backcross generation is then selfed to give pure breeding progeny for the gene(s) being transferred. The resulting plants have essentially all of the morphological and physiological characteristics of inbred maize line NP2222, in addition to the single gene trait(s) transferred to the inbred. Preferably, the resulting plants have all of the morphological and physiological characteristics of inbred maize line NP2222, in addition to the single gene trait(s) transferred to the inbred. The exact backcrossing protocol will depend on the trait being altered to determine an appropriate testing protocol. Although backcrossing methods are simplified when the trait being transferred is a dominant allele, a recessive allele may also be transferred. In this instance it may be necessary to introduce a test of the progeny to determine if the desired trait has been successfully transferred.

Many traits have been identified that are not regularly selected for in the development of a new inbred but that can be improved by backcrossing techniques or genetic transformation. Examples of traits transferred to inbred maize line NP2222 include, but are not limited to, waxy starch, herbicide tolerance, resistance for bacterial, fungal, or viral disease, insect resistance, enhanced nutritional quality, improved performance in an industrial process, altered reproductive capability, such as male sterility or male fertility, yield stability and yield enhancement. Other traits transferred to inbred maize line NP2222 are for the production of commercially valuable enzymes or metabolites in plants of inbred maize line NP2222.

Traits transferred to maize inbred line NP2222 are naturally occurring maize traits, which are preferably introgressed into inbred maize line NP2222 by breeding methods such as backcrossing, or are heterologous transgenes, which are preferably first introduced into a maize line by genetic transformation using genetic engineering and transformation techniques well known in the art, and then introgressed into inbred line NP2222. Alternatively a heterologous trait is directly introduced into inbred maize line NP2222 by genetic transformation. Heterologous, as used herein, means of different natural origin or represents a non-natural state. For example, if a host cell is transformed with a nucleotide sequence derived from another organism, particularly from another species, that nucleotide sequence is heterologous with respect to that host cell and also with respect to descendants of the host cell which carry that gene. Similarly, heterologous refers to a nucleotide sequence derived from and inserted into the same natural, original cell type, but which is present in a non-natural state, e.g. a different copy number, or under the control of different regulatory sequences. A transforming nucleotide sequence may comprise a heterologous coding sequence, or heterologous regulatory sequences. Alternatively, the transforming nucleotide sequence may be completely heterologous or may comprise any possible combination of heterologous and endogenous nucleic acid sequences.

A transgene introgressed into maize inbred line NP2222 typically comprises a nucleotide sequence whose expression is responsible or contributes to the trait under the control of a promoter appropriate for the expression of the nucleotide sequence at the desired time in the desired tissue or part of the plant. Constitutive or inducible promoters

are used. The transgene may also comprise other regulatory elements such as for example translation enhancers or termination signals. In a preferred embodiment, the nucleotide sequence is the coding sequence of a gene and is transcribed and translated into a protein. In another preferred embodiment, the nucleotide sequence encodes an antisense RNA, a sense RNA that is not translated or only partially translated, a t-RNA, a r-RNA or a sn-RNA.

Where more than one trait are introgressed into inbred maize line NP2222, it is preferred that the specific genes are all located at the same genomic locus in the donor, non-recurrent parent, preferably, in the case of transgenes, as part of a single DNA construct integrated into the donor's genome. Alternatively, if the genes are located at different genomic loci in the donor, non-recurrent parent, backcrossing allows to recover all of the morphological and physiological characteristics of inbred maize line NP2222 in addition to the multiple genes in the resulting maize inbred line.

The genes responsible for a specific, single gene trait are generally inherited through the nucleus. Known exceptions are, e.g. the genes for male sterility, some of which are inherited cytoplasmically, but still act as single gene traits. In a preferred embodiment, a heterologous transgene to be transferred to maize inbred line NP2222 is integrated into the nuclear genome of the donor, non-recurrent parent. In another preferred embodiment, a heterologous transgene to be transferred to into maize inbred line NP2222 is integrated into the plastid genome of the donor, non-recurrent parent. In a preferred embodiment, a plastid transgene comprises one gene transcribed from a single promoter or two or more genes transcribed from a single promoter.

In a preferred embodiment, a transgene whose expression results or contributes to a desired trait to be transferred to maize inbred line NP2222 comprises a virus resistance trait such as, for example, a MDMV strain B coat protein gene whose expression confers resistance to mixed infections of maize dwarf mosaic virus and maize chlorotic mottle virus in transgenic maize plants (Murry et al. Biotechnology (1993) 11:1559-64). In another preferred embodiment, a transgene comprises a gene encoding an insecticidal protein, such as, for example, a crystal protein of *Bacillus thuringiensis* or a vegetative insecticidal protein from *Bacillus cereus*, such as VIP3 (see for example Estruch et al. Nat Biotechnol (1997) 15:137-41). In a preferred embodiment, an insecticidal gene

introduced into maize inbred line NP2222 is a Cry1Ab gene or a portion thereof, for example introgressed into maize inbred line NP2222 from a maize line comprising a Bt-11 event as described in U.S. Patent No. 6,114,608, which is incorporated herein by reference, or from a maize line comprising a 176 event as described in Koziel *et al.* (1993) *Biotechnology* 11: 194-200. In yet another preferred embodiment, a transgene introgressed into maize inbred line NP2222 comprises a herbicide tolerance gene. For example, expression of an altered acetohydroxyacid synthase (AHAS) enzyme confers upon plants tolerance to various imidazolinone or sulfonamide herbicides (U.S. Patent No. 4,761,373). In another preferred embodiment, a non-transgenic trait conferring tolerance to imidazolinones is introgressed into maize inbred line NP2222 (e.g. a "IT" or "IR" trait). U.S. Patent No. 4,975,374, incorporated herein by reference, relates to plant cells and plants containing a gene encoding a mutant glutamine synthetase (GS) resistant to inhibition by herbicides that are known to inhibit GS, e.g. phosphinothricin and methionine sulfoximine. Also, expression of a *Streptomyces bar* gene encoding a phosphinothricin acetyl transferase in maize plants results in tolerance to the herbicide phosphinothricin or glufosinate (U.S. Patent No. 5,489,520). U.S. Patent No. 5,013,659, which is incorporated herein by reference, is directed to plants that express a mutant acetolactate synthase (ALS) that renders the plants resistant to inhibition by sulfonylurea herbicides. U.S. Patent No. 5,162,602 discloses plants tolerant to inhibition by cyclohexanedione and aryloxyphenoxypropanoic acid herbicides. The tolerance is conferred by an altered acetyl coenzyme A carboxylase (ACCase). U.S. Patent No. 5,554,798 discloses transgenic glyphosate tolerant maize plants, which tolerance is conferred by an altered 5-enolpyruvyl-3-phosphoshikimate (EPSP) synthase gene. U.S. Patent No. 5,804,425 discloses transgenic glyphosate tolerant maize plants, which tolerance is conferred by an EPSP synthase gene derived from *Agrobacterium tumefaciens* CP-4 strain. Also, tolerance to a protoporphyrinogen oxidase inhibitor is achieved by expression of a tolerant protoporphyrinogen oxidase enzyme in plants (U.S. Patent No. 5,767,373). Another trait transferred to inbred maize line NP2222 confers tolerance to an inhibitor of the enzyme hydroxyphenylpyruvate dioxygenase (HPPD) and transgenes conferring such trait are, for example, described in WO 9638567, WO 9802562, WO 9923886, WO 9925842, WO 9749816, WO 9804685 and WO 9904021.

All issued patents referred to herein are, in their entirety, expressly incorporated herein by reference.

In a preferred embodiment, a transgene transferred to maize inbred line NP2222 comprises a gene conferring tolerance to a herbicide and at least another nucleotide sequence encoding another trait, such as for example, an insecticidal protein. Such combination of single gene traits is for example a Cry1Ab gene and a *bar* gene.

Specific transgenic events introgressed into maize inbred line NP2222 are found at http://www.aphis.usda.gov/bbep/bp/not_reg.html. For example, introgressed from glyphosate tolerant event GA21 (9709901p), glyphosate tolerant/Lepidopteran insect resistant event MON 802 (9631701p), Lepidopteran insect resistant event DBT418 (9629101p), male sterile event MS3 (9522801p), Lepidopteran insect resistant event Bt11 (9519501p), phosphinothricin tolerant event B16 (9514501p), Lepidopteran insect resistant event MON 80100 (9509301p), phosphinothricin tolerant events T14, T25 (9435701p), Lepidopteran insect resistant event 176 (9431901p).

The introgression of a Bt11 event into a maize line, such as maize inbred line NP2222, by backcrossing is exemplified in U.S. Patent No. 6,114,608, and the present invention is directed to methods of introgressing a Bt11 event into maize inbred line NP2222 using for example the markers described in U.S. Pat. No. 6,114,608 and to resulting maize lines.

Direct selection may be applied where the trait acts as a dominant trait. An example of a dominant trait is herbicide tolerance. For this selection process, the progeny of the initial cross are sprayed with the herbicide prior to the backcrossing. The spraying eliminates any plant which does not have the desired herbicide tolerance characteristic, and only those plants that have the herbicide tolerance gene are used in the subsequent backcross. This process is then repeated for the additional backcross generations.

This invention also is directed to methods for producing a maize plant by crossing a first parent maize plant with a second parent maize plant wherein either the first or second parent maize plant is a maize plant of inbred line NP2222 or a maize plant of inbred line NP2222 further comprising one or more single gene traits. Further, both first and second parent maize plants can come from the inbred maize line NP2222 or an inbred maize plant of NP2222 further comprising one or more single gene traits. Thus,

any such methods using the inbred maize line NP2222 or an inbred maize plant of NP2222 further comprising one or more single gene traits are part of this invention: selfing, backcrosses, hybrid production, crosses to populations, and the like. All plants produced using inbred maize line NP2222 or inbred maize plants of NP2222 further comprising one or more single gene traits as a parent are within the scope of this invention. Advantageously, inbred maize line NP2222 or inbred maize plants of NP2222 further comprising one or more single gene traits are used in crosses with other, different, maize inbreds to produce first generation (F1) maize hybrid seeds and plants with superior characteristics.

In a preferred embodiment, seeds of inbred maize line NP2222 or seeds of inbred maize plants of NP2222 further comprising one or more single gene traits are provided as an essentially homogeneous population of inbred corn seeds. Essentially homogeneous populations of inbred seed are those that consist essentially of the particular inbred seed, and are generally purified free from substantial numbers of other seed, so that the inbred seed forms between about 90% and about 100% of the total seed, and preferably, between about 95% and about 100% of the total seed. Most preferably, an essentially homogeneous population of inbred corn seed will contain between about 98.5%, 99%, 99.5% and about 100% of inbred seed, as measured by seed grow outs. The population of inbred corn seeds of the invention is further particularly defined as being essentially free from hybrid seed. The inbred seed population may be separately grown to provide an essentially homogeneous population of plants of inbred maize line NP2222 or inbred maize plants of NP2222 further comprising one or more single gene traits.

As used herein, the term "plant" includes plant cells, plant protoplasts, plant cell tissue cultures from which maize plants can be regenerated, plant calli, plant clumps, and plant cells that are intact in plants or parts of plants, such as embryos, pollen, ovules, flowers, kernels, ears, cobs, leaves, husks, stalks, roots, root tips, anthers, silk, seeds and the like.

Duncan, Williams, Zehr, and Widholm, *Planta* (1985) 165:322-332 reflects that 97% of the plants cultured that produced callus were capable of plant regeneration. Subsequent experiments with both inbreds and hybrids produced 91% regenerable callus that produced plants. In a further study in 1988, Songstad, Duncan & Widholm in *Plant*

Cell Reports (1988), 7:262-265 reports several media additions that enhance regenerability of callus of two inbred lines. Other published reports also indicated that "nontraditional" tissues are capable of producing somatic embryogenesis and plant regeneration. K. P. Rao, et al., Maize Genetics Cooperation Newsletter, 60:64-65 (1986), refers to somatic embryogenesis from glume callus cultures and B. V. Conger, et al., Plant Cell Reports, 6:345-347 (1987) indicates somatic embryogenesis from the tissue cultures of maize leaf segments. Thus, it is clear from the literature that the state of the art is such that these methods of obtaining plants are, and were, "conventional" in the sense that they are routinely used and have a very high rate of success.

Tissue culture procedures of maize are described in Green and Rhodes, "Plant Regeneration in Tissue Culture of Maize," Maize for Biological Research (Plant Molecular Biology Association, Charlottesville, Va. 1982, at 367-372) and in Duncan, et al., "The Production of Callus Capable of Plant Regeneration from Immature Embryos of Numerous Zea mays Genotypes," 165 Planta 322-332 (1985). Thus, another aspect of this invention is to provide cells that upon growth and differentiation produce maize plants having the physiological and morphological characteristics of inbred maize line NP2222. In a preferred embodiment, cells of inbred maize line NP2222 are transformed genetically, for example with one or more genes described above, for example by using a transformation method described in U.S. Pat. No. 6,114,608, and transgenic plants of inbred maize line NP2222 are obtained and used for the production of hybrid maize plants.

Maize is used as human food, livestock feed, and as raw material in industry. The food uses of maize, in addition to human consumption of maize kernels, include both products of dry- and wet-milling industries. The principal products of maize dry milling are grits, meal and flour. The maize wet-milling industry can provide maize starch, maize syrups, and dextrose for food use. Maize oil is recovered from maize germ, which is a by-product of both dry- and wet-milling industries.

Maize, including both grain and non-grain portions of the plant, is also used extensively as livestock feed, primarily for beef cattle, dairy cattle, hogs, and poultry. Industrial uses of maize include production of ethanol, maize starch in the wet-milling industry and maize flour in the dry-milling industry. The industrial applications of maize starch and

flour are based on functional properties, such as viscosity, film formation, adhesive properties, and ability to suspend particles. The maize starch and flour have application in the paper and textile industries. Other industrial uses include applications in adhesives, building materials, foundry binders, laundry starches, explosives, oil-well muds, and other mining applications. Plant parts other than the grain of maize are also used in industry: for example, stalks and husks are made into paper and wallboard and cobs are used for fuel and to make charcoal.

The seed of inbred maize line NP2222 or of inbred maize line NP2222 further comprising one or more single gene traits, the plant produced from the inbred seed, the hybrid maize plant produced from the crossing of the inbred, hybrid seed, and various parts of the hybrid maize plant can be utilized for human food, livestock feed, and as a raw material in industry.

The present invention therefore also discloses an agricultural product comprising a plant of the present invention or derived from a plant of the present invention. The present invention also discloses an industrial product comprising a plant of the present invention or derived from a plant of the present invention. The present invention further discloses methods of producing an agricultural or industrial product comprising planting seeds of the present invention, growing plant from such seeds, harvesting the plants and processing them to obtain an agricultural or industrial product.

DEPOSIT

Applicants have made a deposit of at least 2500 seeds of Inbred Maize Line NP2222 with the American Type Culture Collection (ATCC), Manassas, Virginia, 20110-2209 U.S.A., ATCC Deposit No: PTA-3967. This deposit of the Inbred Maize Line NP2222 will be maintained in the ATCC depository, which is a public depository, for a period of 30 years, or 5 years after the most recent request, or for the effective life of the patent, whichever is longer, and will be replaced if it becomes nonviable during that period. Additionally, Applicants have satisfied all the requirements of 37 C.F.R. §§1.801-1.809, including providing an indication of the viability of the sample. Applicants impose no restrictions on the availability of the deposited material from the ATCC; however, Applicants have no authority to waive any restrictions imposed by law

on the transfer of biological material or its transportation in commerce. Applicants do not waive any infringement of its rights granted under this patent or under the Plant Variety Protection Act (7 USC 2321 et seq.).

5 The foregoing invention has been described in detail by way of illustration and example for purposes of clarity and understanding. However, it will be obvious that certain changes and modifications such as single gene modifications and mutations, somaclonal variants, variant individuals selected from large populations of the plants of the instant inbred and the like may be practiced within the scope of the invention, as limited only by the scope of the appended claims.

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
220